Pediatric Symptom Checklist PSC-17 Description

The PSC-17 is a general mental health screening tool designed to be simple to use in primary care practices, based a longer form instrument known as the PSC-35. It can help primary care providers assess the likelihood of finding any mental health disorder in their patient. The brief and easy to score PSC-17 has fairly good mental health screening characteristics, even when compared with much longer instruments like the CBCL (Child Behavior Checklist by T. Achenbach).

A 2007 study in primary care offices compared use of the PSC-17 to simultaneous use of the CBCL in 269 children aged 8-15, showing reasonably good performance of its three subscales compared to similar subscales on the CBCL. The gold standard here was a K-SADS diagnosis, which is a standardized psychiatric interview diagnosis. These comparison statistics are summarized below, with positive and negative predictive values shown based on different presumed prevalence (5 or 15%) of the disorders. Providers should notice that despite its good performance relative to longer such measures, it is not a foolproof diagnostic aid. For instance the sensitivity for this scale only ranges from 31% to 73% depending on the disorder in this study:

K-SADS Diagnosis	Screen	Sensitivity	Specificity	PPV 5%	PPV 15%	NPV 5%	NPV 15%
ADHD	PSC-17 Attention CBCL Attention	0.58 0.68	0.91 0.90	0.25 0.26	0.53 0.55	0.98 0.98	0.92 0.94
Anxiety	PSC-17 Internalizing CBCL Internalizing	0.52 0.42	0.74 0.88	0.10 0.13	0.26 0.38	0.97 0.97	0.90 0.90
Depression	PSC-17 Internalizing CBCL Internalizing	0.73 0.58	0.74 0.87	0.13 0.19	0.33 0.44	0.98 0.98	0.94 0.92
Externalizing	PSC-17 Externalizing CBCL Externalizing	0.62 0.46	0.89 0.95	0.23 0.33	0.50 0.62	0.98 0.97	0.93 0.91
Any Diagnosis	PSC-17 Total CBCL Total	0.42 0.31	0.86 0.96	0.14 0.29	0.35 0.58	0.97 0.96	0.89 0.89

W Gardner, A Lucas, DJ Kolko, JV Campo "Comparison of the PSC-17 and Alternative Mental Health Screens in an At-Risk Primary Care Sample" JAACAP 46:5, May 2007, 611-618

PSC-17 Scoring:

PSC-17 Internalizing score positive if ≥ 5

PSC-17 Externalizing score positive if ≥ 7

PSC-17 Attention score positive if ≥7

PSC-17 Total score positive if ≥15

"Attention" diagnoses can include: ADHD, ADD

"Internalizing" diagnoses can include: Any anxiety or mood disorder

"Externalizing" diagnoses can include: Conduct disorder, Oppositional Defiant Disorder, adjustment disorder with disturbed conduct or mixed disturbed mood and conduct

Pediatric Symptom Checklist-17 (PSC-17)

	Please tha	_	For Office Use			
	NEVER	SOMETIMES	OFTEN	I	А	Е
Fidgety, unable to sit still						
Feels sad, unhappy						
. Daydreams too much						
. Refuses to share						
Does not understand other people's feelings						
. Feels hopeless						
Has trouble concentrating						
. Fights with other children						
. Is down on him or herself						
D. Blames others for his or her troubles						
. Seems to be having less fun						
2. Does not listen to rules						
3. Acts as if driven by a motor						
1. Teases others						
5. Worries a lot						
6. Takes things that do not belong to him or her						
7. Distracted easily						
(scoring totals)						

Scoring:

- Fill in unshaded box on right with: "Never" = 0, "Sometimes" = 1, "Often" = 2
- Sum the columns.

PSC17 Internalizing score is sum of column I PSC17 Attention score is sum of column A PSC17 Externalizing score is sum of column E PSC-17 Total Score is sum of I, A, and E columns

Suggested Screen Cutoff:

PSC-17 - I ≥ 5

 $PSC-17 - A \ge 7$

PSC-17 - E ≥7

Total Score ≥ 15

Higher Scores can indicate an increased likelihood of a behavioral health disorder being present.

PSC-17 may be freely reproduced.

Created by W Gardner and K Kelleher (1999), and based on PSC by M Jellinek et al. (1988) Formatted by R Hilt, inspired by Columbus Children's Research Institute formatting of PSC-17